

Anti-3PGDH

(Phgdh, 3-phosphoglycerate dehydrogenase)

Product Information:

1) Catalog #:	MSFR100010
Item Name:	3PGDH pAb (Rb) 20ug
Size :	20µg (affinity-purified with antigen polypeptide)
Species :	Rabbit
Product Code :	3PGDH-Rb-Af303
RRID :	AB_2571653
2) Catalog #:	MSFR100020
Item Name:	3PGDH pAb (Rb) 50ug
Size :	50µg (affinity-purified with antigen polypeptide)
Species :	Rabbit
Product Code :	3PGDH-Rb-Af303
RRID :	AB_2571653
3) Catalog #:	MSFR100030
Item Name:	3PGDH pAb (GP) 20ug
Size :	20µg (affinity-purified with antigen polypeptide)
Species :	Guinea pig
Product Code :	3PGDH-GP-Af198
RRID :	AB_2571654
4) Catalog #:	MSFR100040
Item Name:	3PGDH pAb (GP) 50ug
Size :	50µg (affinity-purified with antigen polypeptide)
Species :	Guinea pig
Product Code :	3PGDH-GP-Af198
RRID :	AB_2571654

Formulation : Liquid ; 200 µg/ml in PBS with 0.05% NaN₃.
(affinity-purified with antigen polypeptide)

Storage : Store at cool temp (2-10°C).

The antibody can be stored at 2-10°C. The antibody can be also aliquoted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.

Species : rabbit / guinea pig, polyclonal


Antigen : rat, full sequence (NM031620)

ご注意 : 本商品には0.1%未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD Manufacturer : FRONTIER INSTITUTE CO., LTD
TEL : +81-(0)3-4582-5451 email : nmd-tokuhan@nittobogrp.com

Specificity : Mouse (others not tested)

Immunoblot detects a single band at 57 kDa. This selectively stains astroglia in the central nervous system, Schwann cells in the peripheral nervous system, and distinct tubular segments in the kidney.



Applications : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

Research Use : For research use only, not for use in diagnostic procedures.

Remarks : Rabbit and guinea pig antibodies are similar in the titer and specificity.

Reference : 1) Yamasaki, M., Yamada, K., Furuya, S., Mitoma, J., Hirabayashi, Y., Watanabe, M. (2001) 3-phosphoglycerate dehydrogenase (3PGDH), a key enzyme of L-serine biosynthesis, is preferentially expressed in the radial glia/astrocyte lineage and olfactory ensheathing glia in the mouse brain. **J. Neurosci.** 21:7691-7704.

2) Sakai, K., Shimizu, H., Koike, T., Furuya S., Watanabe, M. (2003) Neutral amino acid transporter ASCT1 is preferentially expressed in L-Ser-synthetic/storing glial cells in the mouse with transient expression in developing capillaries. **J. Neurosci.** 23:550-560.

3) Yamashita, N., Sakai, K., Furuya, S., Watanabe, M. (2003) Selective expression of L-serine synthetic enzyme 3PGDH in Schwann cells, perineuronal glia, and endoneurial fibroblasts along sciatic nerves and its upregulation after crush injury. **Arch. Histol. Cytol.** 66:429-436.

4) Takasaki C, Miura E, Watanabe M: (2007) Segmental and complementary expression of L-serine biosynthetic enzyme 3-phosphoglycerate dehydrogenase and neutral amino acid transporter ASCT1 in the mouse kidney. **Biomed Res** 28(2):61-69

ご注意 : 本商品には 0.1%未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD Manufacturer : FRONTIER INSTITUTE CO., LTD
TEL : +81-(0)3-4582-5451 email : nmd-tokuhan@nittobogrp.com