

Anti-CCK antibody

(cholecystokinin)

Code Number : CCK8-MO-167-2 (clone# 167, RRID : AB_2572276)

: CCK8-MO-1E3-1 (clone# 1E3, RRID : AB_2572277)

Size : 50 µg and 100 µg / See label on vial

(affinity-purified, Protein G)

Formulation : Liquid ;

(CCK8-MO-167-2, 1.0 mg / ml in PBS with 0.03% ProClin)

(CCK8-MO-1E3-1, 1.0 mg / ml in PBS with 0.03% ProClin)

Storage : Store at cool temp. (2-10°C)

The antibody can be stored at 2-10°C. The antibody can be also aliquotted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardrous. No MSDS required.

Species : mouse, monoclonal

Isotype : IgG1



Antigen : synthetic peptide (sulfated CCK-8)

Specificity : mouse, rat (others not tested)



CCK8-MO-167 (clone#167) reacts with the CCK, but does not cross-react with gastrin.

CCK8-MO-1E3 (clone#1E3) cross-react with gastrin.

clone # 167



clone # 1E3



mouse duodenum
paraffin-embedded section

mouse antrum
paraffin-embedded section

ご注意 : 本商品には 0.03 % の ProClin が入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD

Manufacture : FRONTIER INSTITUTE CO., LTD

TEL : +81-(0)3-4582-5451

email : nmd-tokuhan@nittobogrp.com

Applications : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

Research Use : For research use only, not for use in diagnostic procedures.

Remarks : These IHC data were provided by Professor Dr. Toshihiko Iwanaga (Hokkaido University Graduate School of Medicine, Department. of Functional Morphology). Special thanks to his support.

Reference :

ご注意 : 本商品には 0.03 % の ProClin が入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD

Manufacture : FRONTIER INSTITUTE CO., LTD

TEL : +81-(0)3-4582-5451

email : nmd-tokuhan@nittobogrp.com