

immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

Research Use : For research use only, not for use in diagnostic procedures.

Remarks : This stains nearly a half population of striatal medium spiny neurons.

Reference : 1) Narushima, M., Uchigashima, M., Hashimoto, K., Watanabe, M., Kano, M. (2006) Depolarization-induced suppression of inhibition mediated by endocannabinoid at synapses from fast-spiking interneurons to medium spiny neurons in the striatum. **Eur. J. Neurosci.** 24:2246-2252.

2) Uchigashima M, Narushima M, Fukaya M, Katona I, Kano M, Watanabe M: Subcellular arrangement of molecules for 2-arachidonoyl-glycerol-mediated retrograde signaling and its physiological contribution to synaptic modulation in the striatum. **J. Neurosci.**, 27:3663-3676, 2007

ご注意：本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD
TEL : +81-(0)3-4582-5451
email : nmd-tokuhan@nittobogrp.com

Manufacture : FRONTIER INSTITUTE CO., LTD