

## *Anti-GluA1 (GluR1)*

(AMPA-type glutamate receptor subunit-1)

### Product Information:

---

1) Catalog #:	MSFR102270
Item Name:	GluA1 (GluR1)pAb (Rb) 20ug
Size :	20µg (affinity-purified with antigen polypeptide)
Species :	Rabbit
Product Code :	GluA1-Rb-Af690
RRID :	AB_2571752
2) Catalog #:	MSFR102280
Item Name:	GluA1 (GluR1) pAb (Rb) 50ug
Size :	50µg (affinity-purified with antigen polypeptide)
Species :	Rabbit
Product Code :	GluA1-Rb-Af690
RRID :	AB_2571752
3) Catalog #:	MSFR102290
Item Name:	GluA1 (GluR1) pAb (GP) 20ug
Size :	20µg (affinity-purified with antigen polypeptide)
Species :	Guinea pig
Product Code :	GluA1-GP-Af380
RRID :	AB_2571753
4) Catalog #:	MSFR102300
Item Name:	GluA1 (GluR1) pAb (GP) 50ug
Size :	50µg (affinity-purified with antigen polypeptide)
Species :	Guinea pig
Product Code :	GluA1-GP-Af380
RRID :	AB_2571753

---


**Formulation** : Liquid ; 200 µg/ml in PBS with 0.05% NaN<sub>3</sub>.

**Storage** : Store at cool temp. (2-10°C)

The antibody can be stored at 2-10°C. The antibody can be also aliquoted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.

**Species** : rabbit / guinea pig, polyclonal

**Antigen** : mouse GluA1, (X57497)  
841-907 aa, GFCLIPQQSINEAIRTST  
LPRNSGAGASGGSGSGENGRVVSQD  
FPKSMQSIPCMSHSSGMPLGATGL



**Specificity** : mouse (others not tested)

Immunoblot detects a single protein band at 100kDa, with no cross reactivity to other iGluR subunits. Lack of signal in GluA1-KO brain.

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

**Applications** : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

**Research Use** : For research use only, not for use in diagnostic procedures.

**Remarks** : For immunohistochemistry for neuronal iGluRs, users should adopt postembedding immunogold for electron microscopic detection and protease predigestion for light microscopic detection (see the below reference). For glial GluR, these antigen-exposing methods are not necessary (unpublished information).

**Reference** : 1) Watanabe, M., Fukaya, M., Sakimura, K., Manabe, T., Mishina, M., and Inoue, Y. (1998) Selective scarcity of NMDA receptor channel subunits in the stratum lucidum (mossy fiber-recipient layer) of the hippocampal CA3 subfield. **Eur. J. Neurosci.** 10:478-487.  
2) Shimuta, M., Yoshikawa, M., Fukaya, M., Watanabe, M., Takeshima, H., Manabe, T. (2001) Postsynaptic modulation of AMPA receptor-mediated synaptic responses and LTP by the type 3 ryanodine receptor. **Mol. Cell Neurosci.** 17:921-930.  
3) Fukaya, M., Tsujita, M., Yamazaki, M., Kushiya, E., Abe, M., Akashi, K., Natsume, R., Kano, M., Kamiya, H., Watanabe, M. \*, Sakimura, K. Abundant distribution of TARP  $\gamma$ -8 in synaptic and extrasynaptic surface of hippocampal neurons and its major role in AMPA receptor expression on spines and dendrites. **Eur. J. Neurosci.**, 24:2177-2190.

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。