

Anti- GluA2 (GluR2)

(AMPA-type glutamate receptor subunit-2)

Product Information:

1) Catalog #:	MSFR102320
Item Name:	GluA2 (GluR2) pAb (Rb) 20ug
Size :	20µg (affinity-purified with antigen polypeptide)
Species :	Rabbit
Product Code :	GluR2C-Rb-Af1050
RRID :	AB_2571754
2) Catalog #:	MSFR102330
Item Name:	GluA2 (GluR2) pAb (Rb) 50ug
Size :	50µg (affinity-purified with antigen polypeptide)
Species :	Rabbit
Product Code :	GluR2C-Rb-Af1050
RRID :	AB_2571754


Formulation : Liquid ; 200 µg/ml in PBS with 0.05% NaN₃.

Storage : Store at cool temp. (2-10°C)

The antibody can be stored at 2-10°C. The antibody can be also aliquoted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.

Species : rabbit, polyclonal

Antigen : mouse GluR2, C-terminal 847-863 aa
(X57498 ; KVAKNAQNINPSSSQNS)



Specificity : mouse (others not tested)

Immunoblot detects a single protein band at 100kDa, with no cross reactivity to other iGluR subunits, including GluR3.

Applications : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

Research Use : For research use only, not for use in diagnostic procedures.

Remarks : For immunohistochemistry for neuronal iGluRs, users should adopt postembedding immunogold for electron microscopic detection and protease predigestion for light microscopic detection (see the below reference). For glial GluR, these antigen-exposing methods are not necessary (unpublished information).

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Reference : 1) Shimuta, M., Yoshikawa, M., Fukaya, M., Watanabe, M., Takeshima, H., Manabe, T. (2001) Postsynaptic modulation of AMPA receptor-mediated synaptic responses and LTP by the type 3 ryanodine receptor. **Mol. Cell Neurosci.** 17:921-930.
2) Fukaya, M., Tsujita, M., Yamazaki, M., Kushiya, E., Abe, M., Akashi, K., Natsume, R., Kano, M., Kamiya, H., Watanabe, M. *, Sakimura, K. Abundant distribution of TARPs γ -8 in synaptic and extrasynaptic surface of hippocampal neurons and its major role in AMPA receptor expression on spines and dendrites. **Eur. J. Neurosci.**, 24:2177-2190.

ご注意：本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD
TEL : +81-(0)3-4582-5451
email : nmd-tokuhan@nittobogrp.com

Manufacture : FRONTIER INSTITUTE CO., LTD