

Anti-HTT / SERT

(plasmalemmal serotonin transporter)

Code Number : HTT-Rb-Af560 (rabbit, RRID : AB_2571775)
 : HTT-Go-Af970 (goat, RRID : AB_2571776)
 : HTT-GP-Af1400 (guinea pig, RRID : AB_2571777)

Size : 20 µg and 50 µg / See label on vial
(affinity-purified with antigen polypeptide)

Formulation : Liquid ; 200 µg/ml in PBS with 0.05% NaN₃.


Storage : Store at cool temp. (2-10°C)

The antibody can be stored at 2-10°C. The antibody can be also aliquotted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.

Species : rabbit / guinea pig / goat,
polyclonal

Antigen : mouse HTT, 1-77 aa
(AF013604)

Specificity : mouse (others not tested)
Immunoblot detects a single protein band at 67-69 kDa. This selectively stains serotonergic neurons and fibers.



Applications : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

Research Use : For research use only, not for use in diagnostic procedures.

Remarks : Rabbit, guinea pig, and goat antibodies are similar in titer and specificity.

Reference : Somogyi, J., Baude, A., Omori, Y., Shimizu, H., Mestikawy, S.E., Fukaya, M., Ryuichi, S., Watanabe, M., Somogyi, P. (2004) GABAergic basket cells expressing

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

cholecystokinin contain vesicular glutamate transporter type 3 (VGLUT3) in their synaptic terminals in hippocampus and isocortex of the rat. **Eur. J. Neurosci.** 19:552-569

ご注意：本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD
TEL : +81-(0)3-4582-5451
email : nmd-tokuhan@nittobogrp.com

Manufacture : FRONTIER INSTITUTE CO., LTD