

Anti-Homer 1

Product Information:

1)	Catalog #:	MSFR103200
	Item Name:	Homer 1 pAb (Rb) 20ug
	Size :	20µg (affinity-purified with antigen polypeptide)
	Species :	Rabbit
	Product Code :	Homer1-Rb-Af1000
	RRID :	AB_2571774
2)	Catalog #:	MSFR103210
	Item Name:	Homer 1 pAb (Rb) 50ug
	Size :	50µg (affinity-purified with antigen polypeptide)
	Species :	Rabbit
	Product Code :	Homer1-Rb-Af1000
	RRID :	AB_2571774
3)	Catalog #:	MSFR103220
	Item Name:	Homer 1 pAb (Go) 20ug
	Size :	20µg (affinity-purified with antigen polypeptide)
	Species :	Goat
	Product Code :	Homer1-Go-Af1270
	RRID :	AB_2631104
4)	Catalog #:	MSFR103230
	Item Name:	Homer 1 pAb (Go) 50ug
	Size :	50µg (affinity-purified with antigen polypeptide)
	Species :	Goat
	Product Code :	Homer1-Go-Af1270
	RRID :	AB_2631104

Formulation : Liquid ; 200 µg/ml in PBS with 0.05% NaN₃.

Storage : Store at cool temp. (2-10°C)

The antibody can be stored at 2-10°C. The antibody can be also aliquoted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.



Species : rabbit / goat, polyclonal

Antigen : mouse Homer-1, 1-175 aa (NM_147176)

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Specificity : mouse (others not tested)

Immunoblot detects a single protein band at 43-45 kDa.



Applications : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

Research Use : For research use only, not for use in diagnostic procedures.

Remarks : In the Homer family, the first 120 amino acids are highly conserved. This Homer-1 antibody must cross-react Homer-2 and Homer-3, thus working as a pan-Homer antibody. Rabbit and goat Homer-1 antibodies produce similar patterns in immunohistochemistry and immunoblot.

Reference : Nakamura, M., Sato, K., Fukaya, M., Araishi, K., Aiba, A., Kano, M., Watanabe, M. (2004) Signaling complex formation of phospholipase C β 4 with mGluR1 α and IP3R1 at the perisynapse and endoplasmic reticulum in the mouse brain. **Eur. J. Neurosci** 20:2929-2944.
2. Nomura S, Fukaya M, Tsujioka T, Wu D, Watanabe M: Phospholipase C β 3 is distributed in both somatodendritic and axonal compartments and localized around perisynapse and smooth endoplasmic reticulum in mouse Purkinje cell subsets. **Eur. J. Neurosci**. 25:659-672, 2007.
3. Gutierrez-Mecinas M, Polgár E, Watanabe M, Abaira VE, Todd AJ: Immunostaining for Homer reveals the majority of excitatory synapses in the mouse spinal dorsal horn. **Neuroscience**, ;329:171-181, 2016.

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD
TEL : +81-(0)3-4582-5451
email : nmd-tokuhan@nittobogrp.com

Manufacture : FRONTIER INSTITUTE CO., LTD