

## Anti-PLC $\beta$ 4

(phospholipase C $\beta$ 4)

### Product Information:

1) Catalog #:	MSFR105000
Item Name:	PLC $\beta$ 4 pAb (Rb) 20 $\mu$ g
Size :	20 $\mu$ g (affinity-purified with antigen polypeptide)
Species :	Rabbit
Product Code :	PLC $\beta$ 4-Rb-Af81
RRID :	AB_2571831
2) Catalog #:	MSFR105010
Item Name:	PLC $\beta$ 4 pAb (Rb) 50 $\mu$ g
Size :	50 $\mu$ g (affinity-purified with antigen polypeptide)
Species :	Rabbit
Product Code :	PLC $\beta$ 4-Rb-Af81
RRID :	AB_2571831
3) Catalog #:	MSFR107410
Item Name:	PLC $\beta$ 4 pAb (Rb)
Size :	20 $\mu$ g (affinity-purified with antigen polypeptide)
Species :	Rabbit
Product Code :	PLC $\beta$ 4-Rb-Af290
RRID :	AB_2864357
4) Catalog #:	MSFR107420
Item Name:	PLC $\beta$ 4 pAb (Rb)
Size :	50 $\mu$ g (affinity-purified with antigen polypeptide)
Species :	Rabbit
Product Code :	PLC $\beta$ 4-Rb-Af290
RRID :	AB_2864357

**Formulation** : Liquid ; 81  $\mu$ g/ml in PBS with 0.05% NaN<sub>3</sub>.

(affinity-purified with antigen polypeptide)

**Storage** : Store at cool temp (2-10°C).


The antibody can be stored at 2-10°C. The antibody can be also aliquoted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.

**Species** : rabbit, polyclonal

**Antigen** : mouse PLC $\beta$ 4, 15-74 aa  
(AF022804),

**Specificity** : mouse (others not tested)

Immunoblot detects a single band at 134 kDa. The specificity was verified by blank immunostaining in PLC $\beta$ 4-KO brains.



ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

**Applications** : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs. In use of cryosections and microslicer sections, pretreatment of dipping sections in 1:3 methanol, 3:1 methanol, and pure methanol for 3 min each will greatly enhance immunohistochemical signals for PLC $\beta$ .

**Research Use** : For research use only, not for use in diagnostic procedures.

**Remarks** : Avoid freeze-drying of this antibody.

**Reference** : 1) Nakamura, M., Sato, K., Fukaya, M., Araishi, K., Aiba, A., Kano, M., Watanabe, M. (2004) Signaling complex formation of phospholipase C $\beta$ 4 with mGluR1 $\alpha$  and IP3R1 at the perisynapse and endoplasmic reticulum in the mouse brain. **Eur. J. Neurosci** 20:2929-2944.  
2) Yoshida, T., Fukaya, M., Uchigashima, M., Kamiya, H., Kano, M., Watanabe, M. (2006) Localization of diacylglycerol lipase- $\alpha$  around postsynaptic spine suggests close proximity between production site of an endocannabinoid, 2-arachidonoyl-glycerol, and presynaptic cannabinoid CB1 receptor. **J. Neurosci.** 26: 4740-4751.

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD  
TEL : +81-(0)3-4582-5451  
email : nmd-tokuhan@nittobogrp.com

Manufacture : FRONTIER INSTITUTE CO., LTD