

## *Anti-SgII*

*(secretogranin II, Chromogranin C)*

### Product Information:

---

1)	<b>Catalog #:</b>	MSFR105440
	<b>Item Name:</b>	SgII (Chromogranin C) pAb (Rb) 20ug
	<b>Size :</b>	20µg (affinity-purified with antigen polypeptide)
	<b>Species :</b>	Rabbit
	<b>Product Code :</b>	SgII-Rb-Af630
	<b>RRID :</b>	AB_2571837
2)	<b>Catalog #:</b>	MSFR105450
	<b>Item Name:</b>	SgII (Chromogranin C) pAb (Rb) 50ug
	<b>Size :</b>	50µg (affinity-purified with antigen polypeptide)
	<b>Species :</b>	Rabbit
	<b>Product Code :</b>	SgII-Rb-Af630
	<b>RRID :</b>	AB_2571837
3)	<b>Catalog #:</b>	MSFR105460
	<b>Item Name:</b>	SgII (sChromogranin C) pAb (GP) 20ug
	<b>Size :</b>	20µg (affinity-purified with antigen polypeptide)
	<b>Species :</b>	Guinea pig
	<b>Product Code :</b>	SgII-GP-Af750
	<b>RRID :</b>	AB_2571838
4)	<b>Catalog #:</b>	MSFR105470
	<b>Item Name:</b>	SgII (Chromogranin C) pAb (GP) 50ug
	<b>Size :</b>	50µg (affinity-purified with antigen polypeptide)
	<b>Species :</b>	Guinea pig
	<b>Product Code :</b>	SgII-GP-Af750
	<b>RRID :</b>	AB_2571838

---

**Formulation :** Liquid ; 200 µg/ml in PBS with 0.05% NaN<sub>3</sub>.

**Storage :** Store at cool temp (2-10°C).

The antibody can be stored at 2-10°C. The antibody can be also aliquoted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.


**Species :** rabbit / guinea pig, polyclonal

**Antigen :** mouse SgII, 532-577aa (AAH48249)

ご注意：本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

**Specificity** : mouse (others not tested)

Immunoblot detects protein bands at 84 and 86 kDa.



**Applications** : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

**Research Use** : For research use only, not for use in diagnostic procedures.

**Remarks** : If you have no preference for species, rabbit SgII antibody is better to choose, although the specificity is the same.

**Reference** : Miyazaki T, Yamasaki M, Uchigashima M, Matsushima A, Watanabe M (2011) Cellular expression and subcellular localization of secretogranin II in the mouse hippocampus and cerebellum. **Eur J Neurosci.** 33:82-94

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD  
TEL : +81-(0)3-4582-5451  
email : nmd-tokuhan@nittobogrp.com

Manufacture : FRONTIER INSTITUTE CO., LTD