

Anti-Synapsin-I

Product Information:

1)	Catalog #:	MSFR105580
	Item Name:	Synapsin-Ia pAb (Rb) 20ug
	Size :	20µg (affinity-purified with antigen polypeptide)
	Species :	Rabbit
	Product Code :	Synapsin-Ia-Rb-Af670
	RRID :	AB_2636934
2)	Catalog #:	MSFR105590
	Item Name:	Synapsin-Ia pAb (Rb) 50ug
	Size :	50µg (affinity-purified with antigen polypeptide)
	Species :	Rabbit
	Product Code :	Synapsin-Ia-Rb-Af670
	RRID :	AB_2636934
3)	Catalog #:	MSFR105600
	Item Name:	Synapsin-Ib pAb (Rb) 20ug
	Size :	20µg (affinity-purified with antigen polypeptide)
	Species :	Rabbit
	Product Code :	Synapsin-Ib-Rb-Af890
	RRID :	AB_2636935
4)	Catalog #:	MSFR105610
	Item Name:	Synapsin-Ib pAb (Rb) 50ug
	Size :	50µg (affinity-purified with antigen polypeptide)
	Species :	Rabbit
	Product Code :	Synapsin-Ib-Rb-Af890
	RRID :	AB_2636935
5)	Catalog #:	MSFR105620
	Item Name:	Synapsin-Ib pAb (GP) 20ug
	Size :	20µg (affinity-purified with antigen polypeptide)
	Species :	Guinea pig
	Product Code :	Synapsin-Ib-GP-Af710
	RRID :	AB_2636936
6)	Catalog #:	MSFR105630
	Item Name:	Synapsin-Ib pAb (GP) 50ug
	Size :	50µg (affinity-purified with antigen polypeptide)
	Species :	Guinea pig
	Product Code :	Synapsin-Ib-GP-Af710
	RRID :	AB_2636936

Formulation : Liquid ; 200 µg/ml in PBS with 0.05% NaN₃.

Storage : Store at cool temp (2-10°C).


The antibody can be stored at 2-10°C. The antibody can be also aliquoted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.

Species : rabbit / guinea pig, polyclonal

Antigen : mouse synapsin-Ia (686-706aa, NM_013680 ; DEVKAETIRSLRKSFASLFSD) and

ご注意：本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

mouse synapsin-Ib (660-670aa, NM_001110780 ; QLKASPSQAQP). These C-terminal sequences are unique to Ia and Ib splice variants of synapsin-I, respectively. Due to high sequence similarity, however, the antibody directed against synapsin-Ia 686-706aa should also recognize the C-termini of synapsin IIa (DEAKAETIRSLRKSFASLFSD) and III (DEAKAETIRNLRKSFASLFSD). In contrast, the C-terminus used to raise Synapsin-Ib antibody is unique among five synapsin isoforms.



Specificity : In immunoblot, synapsin-Ia antibody detects three protein bands at 60-75 kDa (presumably representing synapsin-Ia, IIa, and III), while synapsin-Ib antibody detect a single protein band at at 75 kDa. When the same blotted membrane is successively immunoreacted with Ia and Ib antibodies, the size of detected bands at around 75 kDa is slightly different (left). Preabsorption of Ia antibody with Ia antigen peptide (686-706aa), but not Ib (660-670aa), abolishes immunosignals (right upper). Likewise, preabsorption of Ib antibody with Ib peptide, but not Ia, abolishes immunosignals (right lower), indicating no cross-reactivity to the other isoforms. In immunohistochemistry, both of synapsin-Ia and -Ib antibodies selectively stains nerve terminals in similar patterns.

Applications : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

Research Use : For research use only, not for use in diagnostic procedures.

Remarks :

Reference :

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。