

## *Anti-TARP $\gamma$ 3*

*(transmembrane AMPA receptor-interacting protein  $\gamma$  3)*

**Code Number** : TARPg3-GP-Af400 (guinea pig, RRID : AB\_2687433)

**Size** : 20  $\mu$ g and 50  $\mu$ g / See label on vial  
(affinity-purified with antigen polypeptide)

**Formulation** : Liquid ; 200  $\mu$ g/ml in PBS with 0.05% NaN<sub>3</sub>.

**Storage** : Store at cool temp (2-10°C).


The antibody can be stored at 2-10°C. The antibody can be also aliquotted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.

**Species** : guinea pig, polyclonal

**Antigen** : mouse, 290-310 aa (FHNSTPKKEFKESLHNNPANR, NM\_019430)

**Specificity** : mouse (others not tested)

This antibody strongly stains cerebellar cortex of wild-type mice, but not its knockout mice.



ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

**Applications** : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

**Research Use** : For research use only, not for use in diagnostic procedures.

**Remarks** : To detect  $\gamma$ -3 at postsynaptic site, antigen retrieval methods need to be adopted, such as pepsin pretreatment of sections or postembedding immunogold microscopy.

**Reference** : 1) Yamasaki M, Fukaya M, Yamazaki M, Azechi H, Natsume R, Abe M, Sakimura K, Watanabe M: TARP  $\gamma$ -2 and  $\gamma$ -8 differentially control biased AMPAR density across Schaffer collateral/commissural synapses in the hippocampal CA1. **J Neurosci**, 36:4296-4312, 2016.

ご注意：本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD  
TEL : +81-(0)3-4582-5451  
email : nmd-tokuhan@nittobogrp.com

Manufacture : FRONTIER INSTITUTE CO., LTD