

Anti-TARP γ 8

(transmembrane AMPA receptor-interacting protein γ 8)

Product Information:

1)	Catalog #:	MSFR105820
	Item Name:	TARP γ 8 pAb (Rb) 20ug
	Size :	20 μ g (affinity-purified with antigen polypeptide)
	Species :	Rabbit
	Product Code :	TARPg8-Rb-Af1000
	RRID :	AB_2571846
2)	Catalog #:	MSFR105830
	Item Name:	TARP γ 8 pAb (Rb) 50ug
	Size :	50 μ g (affinity-purified with antigen polypeptide)
	Species :	Rabbit
	Product Code :	TARPg8-Rb-Af1000
	RRID :	AB_2571846
3)	Catalog #:	MSFR105840
	Item Name:	TARP γ 8 pAb (GP) 20ug
	Size :	20 μ g (affinity-purified with antigen polypeptide)
	Species :	Guinea pig
	Product Code :	TARPg8-GP-Af1000
	RRID :	AB_2572272
4)	Catalog #:	MSFR105850
	Item Name:	TARP γ 8 pAb (GP) 50ug
	Size :	50 μ g (affinity-purified with antigen polypeptide)
	Species :	Guinea pig
	Product Code :	TARPg8-GP-Af1000
	RRID :	AB_2572272

Formulation : Liquid ; 200 μ g/ml in PBS with 0.05% NaN₃.

Storage : Store at cool temp (2-10°C).

The antibody can be stored at 2-10°C. The antibody can be also aliquoted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.

Species : rabbit / guinea pig, polyclonal



Antigen : mouse, 362-421aa

(LTLHNAFPKEAASGVTVTVTGPPAAPAPAPAPPAPAPGTLKSKEAAASNTNTLNRKTT, AF361350), GST fusion (AF361350)

ご注意：本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Specificity : mouse (others not tested)

This antibody stains telencephalic regions of wild-type mice, but not its knockout mice.



Applications : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

Research Use : For research use only, not for use in diagnostic procedures.

Remarks : Rabbit antibody is higher in specificity and titer than guinea pig one.

Reference : Fukaya, M., Tsujita, M., Yamazaki, M., Kushiya, E., Abe, M., Akashi, K., Natsume, R., Kano, M., Kamiya, H., Watanabe, M., Sakimura, K. Abundant distribution of TARP γ -8 in synaptic and extrasynaptic surface of hippocampal neurons and its major role in AMPA receptor expression on spines and dendrites. **Eur. J. Neurosci.**, 24:2177-2190, 2006.
2) Yamasaki M, Fukaya M, Yamazaki M, Azechi H, Natsume R, Abe M, Sakimura K, Watanabe M: TARP γ -2 and γ -8 differentially control biased AMPAR density across Schaffer collateral/commissural synapses in the hippocampal CA1. **J Neurosci**, 36:4296-4312, 2016.

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD
TEL : +81-(0)3-4582-5451
email : nmd-tokuhan@nittobogrp.com

Manufacture : FRONTIER INSTITUTE CO., LTD