

Anti-VGluT3
(vesicular glutamate transporte-3)

Product Information:

1) Catalog #:	MSFR106330
Item Name:	VGluT3 pAb (Go) 20ug
Size :	20µg (affinity-purified with antigen polypeptide)
Species :	Goat
Product Code :	VGluT3-Go-Af870
RRID :	AB_2571854
2) Catalog #:	MSFR106340
Item Name:	VGluT3 pAb (Go) 50ug
Size :	50µg (affinity-purified with antigen polypeptide)
Species :	Goat
Product Code :	VGluT3-Go-Af870
RRID :	AB_2571854
3) Catalog #:	MSFR106350
Item Name:	VGluT3 (rat C-term) pAb (GP) 20ug
Size :	20µg (affinity-purified with antigen polypeptide)
Species :	Guinea pig
Product Code :	VGluT3-GP-Af300
RRID :	AB_2571855
4) Catalog #:	MSFR106360
Item Name:	VGluT3 (rat C-term) pAb (GP) 50ug
Size :	50µg (affinity-purified with antigen polypeptide)
Species :	Guinea pig
Product Code :	VGluT3-GP-Af300
RRID :	AB_2571855
5) Catalog #:	MSFR106370
Item Name:	VGluT3 pAb (GP) 20ug
Size :	20µg (affinity-purified with antigen polypeptide)
Species :	Guinea pig
Product Code :	VGluT3-GP-Af920
RRID :	AB_2571856
6) Catalog #:	MSFR106380
Item Name:	VGluT3 pAb (GP) 50ug
Size :	50µg (affinity-purified with antigen polypeptide)
Species :	Guinea pig
Product Code :	VGluT3-GP-Af920
RRID :	AB_2571856

Formulation : Liquid ; 200 µg/ml in PBS with 0.05% NaN₃.
(affinity-purified with antigen polypeptide)

Storage : Store at cool temp (2-10°C).

The antibody can be stored at 2-10°C. The antibody can be also aliquoted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardous. No MSDS required.

Species : guinea pig / goat, polyclonal

ご注意 : 本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Antigen : mouse VGluT3, 522-588 aa


(MSYGATTQNCEVQKTEWRQQRESAFDGEEPLSYQAEGDFSETS, (AF510321)

VGluT3-GP-Af300 is against the corresponding region of rat VGluT3 (MSYGATTQNCEVQKT
DRRQQRESAFEGEEPLSYQNEEDFSETS, NM153725).

Users should be aware of frequent amino acid substitution between mouse and rat sequences.

Specificity : mouse (others not tested)

Immunoblot detects a single protein band at 60-62 kDa. This stains distinct neuronal populations, which have not been classified as glutamatergic neurons.



Applications : In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

Research Use : For research use only, not for use in diagnostic procedures.

Remarks : Guinea pig and goat antibodies are similar in the titer and specificity.

Reference : Somogyi, J., Baude, A., Omori, Y., Shimizu, H., Mestikawy, S.E., Fukaya, M., Ryuichi, S., Watanabe, M., Somogyi, P. (2004) GABAergic basket cells expressing cholecystokinin contain vesicular glutamate transporter type 3 (VGLUT3) in their synaptic terminals in hippocampus and isocortex of the rat. **Eur. J. Neurosci.** 19:552-569.

ご注意：本商品には 0.1 % 未満のアジ化ナトリウムが入っています。誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor : NITTOBO MEDICAL CO., LTD
TEL : +81-(0)3-4582-5451
email : nmd-tokuhan@nittobogrp.com

Manufacture : FRONTIER INSTITUTE CO., LTD