

## Anti-mGluR5

(metabtropic glutamate receptor-5)

Code Number: mGluR5-Rb-Af300 (rabbit, RRID: AB\_2571802)

: mGluR5-Go-Af540 (goat, RRID: AB\_2571803)

Size: 20 μg and 50 μg / See label on vial

(affinity-purified with antigen polypeptide)

Formulation : Liquid ; 200  $\mu$ g/ml in PBS with 0.05% NaN<sub>3</sub>.

**Storage**: Store at cool temp (2-10°C).

The antibody can be stored at 2-10°C. The antibody can be also aliquotted and stored at -80°C for long-term storage. Avoid repeated freeze-thawing. Non-hazardrous. No MSDS required.

**Species**: rabbit / guinea pig / goat, polyclonal

**Antigen**: mouse mGluR5, C-terminal

1144-1171 aa (D10891)

mGluR5

Specificity: mouse (others not tested)

Immunoblot detects a single protein band at 160 kDa with no cross reactivity to mGluR1. The specificity was verified by blank staining in mGluR5-KO brain.

**Applications**: In general, affinity-purified antibody is used at around 1 microgram/ml for immunoblot and immunohistochemistry. The most appropriate concentration should be determined by users, because it depends on contents in given cells, tissues and organs.

Research Use: For research use only, not for use in diagnostic procedures.

**Remarks**: Rabbit, guinea pig, and goat antibodies are similar in titer and specificity.

**Reference**: 1) Tanaka, J., Nakagawa, S., Yamasaki, M., Fukaya, M., Iwanaga, T., Sakimura, K., Kano, M., Simon, M.I., Inoue, Y., Watanabe, M. (2000) Gq protein α subunits Gαq and Gα11 are localized at postsynaptic extra-junctional membrane of cereballar Purkinje cells and hippocampal pyramidal cells. **Eur. J. Neurosci**. 12:781-792.

ご注意:本商品には 0.1 % 未満のアジ化ナトリウムが入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor: NITTOBO MEDICAL CO., LTD Manufacture: FRONTIER INSTITUTE CO., LTD

TEL: +81-(0)3-4582-5451

email: nmd-tokuhan@nittobogrp.com



2) Uchigashima M, Narushima M, Fukaya M, Katona I, Kano M, Watanabe M: Subcellular arrangement of molecules for 2-arachidonoyl-glycerol-mediated retrograde signaling and its physiological contribution to synaptic modulation in the striatum. **J. Neurosci**. in press.

ご注意:本商品には 0.1% 未満のアジ化ナトリウムが入っています。 誤って目や口に入ったり、皮膚に付着した場合は大量の水で洗い流してください。

Distributor: NITTOBO MEDICAL CO., LTD Manufacture: FRONTIER INSTITUTE CO., LTD

TEL: +81-(0)3-4582-5451

email: nmd-tokuhan@nittobogrp.com